

CP-VIOLATING ASYMMETRIES IN $K^+ \rightarrow 3\pi$ IN THE STANDARD MODEL

ELVIRA GÁMIZ and JOAQUIM PRADES

*CAFPE and Departamento de Física Teórica y del Cosmos
Universidad de Granada, Campus de Fuente Nueva, E-18002 Granada, Spain*

IGNAZIO SCIMEMI

Institute of Theoretical Physics, University of Bern, Sidlerstr. 5, CH-3012 Bern, Switzerland

Abstract

We update the CP-violating asymmetries in $K^+ \rightarrow 3\pi$. In particular we study Δg_C and Δg_N –the asymmetries in the slope g . We emphasize its complementarity to the direct CP-violation parameter ε'_K and the large sensitivity to the size of the imaginary part of the octet coupling $\text{Im } G_8$ at lowest order in CHPT. We also give the prospects of a full calculation at next-to-leading order of this and other CP-violating asymmetries which will be presented elsewhere.

We have calculated¹

$$\begin{aligned} K_2(k) &\rightarrow \pi^0(p_1)\pi^0(p_2)\pi^0(p_3), \\ K_2(k) &\rightarrow \pi^+(p_1)\pi^-(p_2)\pi^0(p_3), \\ K_1(k) &\rightarrow \pi^+(p_1)\pi^-(p_2)\pi^0(p_3), \\ K^+(k) &\rightarrow \pi^0(p_1)\pi^0(p_2)\pi^+(p_3), \\ K^+(k) &\rightarrow \pi^+(p_1)\pi^+(p_2)\pi^-(p_3), \end{aligned} \tag{1}$$

as well as their CP-conjugated decays at one-loop in Chiral Perturbation Theory (CHPT) 2,3,4 –see⁵ for reviews. Above, the momentum of each particle is denoted between brackets and

$$K_{1(2)} = \frac{K^0 - (+)\overline{K}^0}{\sqrt{2}}. \tag{2}$$

The calculation has been done in the limit of equal up and down light quark masses and we have included one-loop electromagnetic corrections for the dominant octet part. We have not included photon loops.

CHPT is the effective field theory of the Standard Model at energies below one GeV . The lowest order $\text{SU}(3) \times \text{SU}(3)$ chiral Lagrangian describing $|\Delta S| = 1$ transitions is

$$\begin{aligned} \mathcal{L}_{|\Delta S|=1}^{(2)} &= C F_0^6 e^2 G_E \text{tr} (\Delta_{32} u^\dagger Q u) + C F_0^4 [G_8 \text{tr} (\Delta_{32} u_\mu u^\mu) + G'_8 \text{tr} (\Delta_{32} \chi_+)] \\ &+ G_{27} t^{ij,kl} \text{tr} (\Delta_{ij} u_\mu) \text{tr} (\Delta_{kl} u^\mu) + \text{h.c.}, \end{aligned} \tag{3}$$

with

$$C = -\frac{3}{5} \frac{G_F}{\sqrt{2}} V_{ud} V_{us}^* \simeq -1.07 \cdot 10^{-6} \text{ GeV}^{-2}. \quad (4)$$

$F_0 = (87 \pm 6) \text{ MeV}$ is the chiral limit value of the pion decay constant $f_\pi = (92.4 \pm 0.4) \text{ MeV}$,

$$\begin{aligned} u_\mu &\equiv iu^\dagger(D_\mu U)u^\dagger = u_\mu^\dagger, \\ \Delta_{ij} &= u\lambda_{ij}u^\dagger, \\ (\lambda_{ij})_{ab} &\equiv \delta_{ia}\delta_{jb}, \\ \chi_{+(-)} &= u^\dagger\chi u^\dagger + (-)u\chi^\dagger u, \end{aligned} \quad (5)$$

$\chi = \text{diag}(m_u, m_d, m_s)$, a 3×3 matrix collecting the light quark masses, and $U \equiv u^2 = \exp(i\sqrt{2}\Phi/F_0)$ is the exponential representation incorporating the octet of light pseudo-scalar mesons in the SU(3) matrix Φ ,

$$\Phi \equiv \begin{pmatrix} \frac{\pi^0}{\sqrt{2}} + \frac{\eta_8}{\sqrt{6}} & \pi^+ & K^+ \\ \pi^- & -\frac{\pi^0}{\sqrt{2}} + \frac{\eta_8}{\sqrt{6}} & K^0 \\ K^- & \bar{K}^0 & -2\frac{\eta_8}{\sqrt{6}} \end{pmatrix}.$$

The non-zero components of the $\text{SU}(3) \times \text{SU}(3)$ tensor $t^{ij,kl}$ are

$$\begin{aligned} t^{21,13} = t^{13,21} &= \frac{1}{3}, & t^{22,23} = t^{23,22} &= -\frac{1}{6}, \\ t^{23,33} = t^{33,23} &= -\frac{1}{6}, & t^{23,11} = t^{11,23} &= \frac{1}{3}, \end{aligned} \quad (6)$$

and $Q = \text{diag}(2/3, -1/3, -1/3)$ is a 3×3 matrix which collects the electric charge of the three light quark flavors. The Lagrangian needed to describe $K \rightarrow 3\pi$ at next-to-leading order can be found in^{3,4,6,7,8,9}.

There is a long history of work concerning $K \rightarrow 3\pi$. First, using current algebra and tree-level chiral Lagrangians¹⁰. The one-loop calculation was done in¹¹, unfortunately the analytical full results were not available. Recently there has appeared the first full published result in¹². We have confirmed this result redoing the one-loop calculation¹.

Using the $K \rightarrow 3\pi$ one-loop results and the ones in¹³ for $K \rightarrow \pi\pi$, a fit to all measured CP-conserving amplitudes and decay rates of those two transitions was done in¹² updating the results in¹¹. The result found there for the ratio of the isospin definite [0 and 2] $K \rightarrow \pi\pi$ amplitudes to all orders in CHPT is

$$\frac{A_0[K \rightarrow \pi\pi]}{A_2[K \rightarrow \pi\pi]} = 21.8, \quad (7)$$

giving the infamous $\Delta I = 1/2$ rule and

$$\left[\frac{A_0[K \rightarrow \pi\pi]}{A_2[K \rightarrow \pi\pi]} \right]^{(2)} = 17.8, \quad (8)$$

to lowest CHPT order p^2 . I.e., Final State Interactions (FSI) and the rest of higher order corrections account for 22 % of the $\Delta I = 1/2$ rule¹². Yet most of this enhancement appears at lowest CHPT order ! The last result is equivalent to the values of the lowest order coupling constants

$$\text{Re } G_8 = 5.5 \pm 0.5 \quad \text{and} \quad G_{27} = 0.39 \pm 0.05. \quad (9)$$

In this normalization, $G_8 = G_{27} = 1$ at large N_c .

CP-violating observables in $K \rightarrow 3\pi$ has attracted also a lot of work since long time ago 14,15,16,17,18,19,21,22 and references therein.

In this talk, we update on the CP-violating asymmetries in the slope g defined as

$$\frac{|A_{K \rightarrow 3\pi}(s_1, s_2, s_3)|^2}{|A_{K \rightarrow 3\pi}(s_0, s_0, s_0)|^2} = 1 + g y + \mathcal{O}(x^2, y^2), \quad (10)$$

with $s_i \equiv (k - p_i)^2$, $3s_0 \equiv m_K^2 + m_{\pi(1)}^2 + m_{\pi(2)}^2 + m_{\pi(3)}^2$ and the Dalitz variables

$$x \equiv \frac{s_1 - s_2}{m_{\pi^+}^2}, \quad y \equiv \frac{s_3 - s_0}{m_{\pi^+}^2}. \quad (11)$$

The two asymmetries that we will be discussing here are

$$\Delta g_C \equiv \frac{g[K^+ \rightarrow \pi^+ \pi^+ \pi^-] - g[K^- \rightarrow \pi^- \pi^- \pi^+]}{g[K^+ \rightarrow \pi^+ \pi^+ \pi^-] + g[K^- \rightarrow \pi^- \pi^- \pi^+]} \quad (12)$$

and

$$\Delta g_N \equiv \frac{g[K^+ \rightarrow \pi^0 \pi^0 \pi^+] - g[K^- \rightarrow \pi^0 \pi^0 \pi^-]}{g[K^+ \rightarrow \pi^0 \pi^0 \pi^+] + g[K^- \rightarrow \pi^0 \pi^0 \pi^-]}. \quad (13)$$

They have been discussed in the literature before finding conflicting results

$$|\Delta g_C| \simeq |\Delta g_N| \simeq 11.0 \times 10^{-4}, \quad (14)$$

in 18,

$$|\Delta g_C| \simeq 0.07 \times 10^{-4}, \quad (15)$$

in 19, and

$$|\Delta g_C| \simeq 0.016 \times 10^{-4}, \quad (16)$$

in 20. The first result was claimed to be at one-loop, however they did not use CHPT fully at one-loop. The last two results are at lowest order p^2 , but they made assumptions to get those results too. Among them, they assumed the exact dominance of the gluonic penguin and neglected the NLO chiral corrections in the result of ε'_K to extract the contribution of the gluonic penguin to the asymmetries. They also made some estimate of the NLO based on strong assumptions which could increase their LO result up to one order of magnitude.

Recently NA48 has announced the possibility of measuring the asymmetry Δg_C with a sensitivity of the order of 10^{-4} and maybe Δg_N , see for instance²³. It is therefore mandatory to have an update of these predictions at NLO in CHPT.

We present here the CHPT leading order results for those asymmetries. The complete NLO analysis including the dominant contributions to the imaginary part at the NLO non-trivial order, i.e. at order p^6 , will be ready soon¹. At LO, the CP-violating asymmetry in the slopes $\Delta g_{C(N)}$ can be written to a very good approximation as

$$\Delta g_{C(N)} \simeq \frac{m_K^2}{F_0^2} A_{C(N)} \text{Im } G_8 + B_{C(N)} \text{Im } (e^2 G_E), \quad (17)$$

with

$$A_{C(N)} = f_{C(N)}(\text{Re } G_8, G_{27}, m_K^2, m_\pi^2) \quad \text{and} \quad B_{C(N)} = h_{C(N)}(\text{Re } G_8, G_{27}, m_K^2, m_\pi^2). \quad (18)$$

The analytic formulas can be found in¹.

Numerically, using the results in (9) for $\text{Re } G_8$ and G_{27} ,

$$\begin{aligned} \Delta g_C &\simeq 0.015 \text{Im } G_8 - 0.0014 \text{Im } (e^2 G_E), \\ \Delta g_N &\simeq -0.0065 \text{Im } G_8 - 0.0008 \text{Im } (e^2 G_E). \end{aligned} \quad (19)$$

Let us see what we do know about the couplings $\text{Im } G_8$ and $\text{Im } G_E$. At large N_c , the contributions to $\text{Im } G_8$ and $\text{Im } G_E$ are factorizable and there is no scheme nor matching scale

dependence. The unfactorizable topologies bring unrelated dynamics, so that we cannot give any uncertainty to the large N_c result. We get

$$\begin{aligned}\text{Im } G_8 \Big|_{N_c} &= (2.0 \pm ?) \text{Im } \tau, \\ \text{Im } (e^2 G_E) \Big|_{N_c} &= -(4.3 \pm ?) \text{Im } \tau.\end{aligned}\quad (20)$$

In the Standard Model

$$\text{Im } \tau \equiv - \left| \frac{V_{td} V_{ts}^*}{V_{ud} V_{us}^*} \right| \simeq -6.05 \times 10^{-4}, \quad (21)$$

and we used here²⁴

$$\langle \bar{q}q \rangle_{\overline{\text{MS}}}(2\text{GeV}) = -(0.018 \pm 0.004) \text{GeV}^3, \quad (22)$$

which agrees with most recent results for the quark condensate and light quark masses from QCD sum rules and light quark masses from lattice QCD, see¹ for references. At this order,

$$\Delta g_C \Big|_{N_c}^{LO} \simeq -0.22 \times 10^{-4} \quad \text{and} \quad \Delta g_N \Big|_{N_c}^{LO} \simeq 0.57 \times 10^{-5}. \quad (23)$$

There have been recently advances on going beyond the leading order in $1/N_c$ in both couplings, $\text{Im } G_8$ and $\text{Im } G_E$.

In^{25,26,27}, there are recent model independent calculations of $\text{Im } G_E$. The results there are valid to all orders in $1/N_c$ and NLO in α_S . They are obtained using hadronic tau data collected by the ALEPH and OPAL Collaborations at LEP. The agreement is quite good between them and their result can be summarized in

$$\text{Im } (e^2 G_E) = -(5.9 \pm 1.3) \text{Im } \tau, \quad (24)$$

where the central value is an average and the error is the smallest one. In²⁸ it was used a Minimal Hadronic Approximation to large N_c to calculate $\text{Im } G_E$, they get

$$\text{Im } (e^2 G_E) = -(9.9 \pm 3.0) \text{Im } \tau, \quad (25)$$

which is also in agreement though somewhat larger. There are also lattice results for $\text{Im } G_E$ using domain-wall fermions²⁹ and Wilson fermions³⁰. All of them made the chiral limit extrapolations, their results are in agreement between them and the average gives

$$\text{Im } (e^2 G_E) = -(4.7 \pm 0.4) \text{Im } \tau. \quad (26)$$

There are also results about $\text{Im } G_8$ at NLO in $1/N_c$. In³¹, the authors did a calculation using a hadronic model which reproduced the $\Delta I = 1/2$ rule through very large Q_2 penguin-like contributions³²

$$\text{Re } G_8 = 5.9 \pm 1.5, \quad \text{and} \quad G_{27} = 0.33 \pm 0.10, \quad (27)$$

in very good agreement with the experimental results in (9). The result found there was at NLO in $1/N_c$

$$\text{Im } G_8 = (5.1 \pm 1.2) \text{Im } \tau. \quad (28)$$

The hadronic model used there had however some drawbacks³³ which will be eliminated and the work eventually updated following the lines in³⁴.

Very recently, using a Minimal Hadronic Approximation to large N_c , the authors of³⁵ found qualitatively similar results. I.e. enhancement toward the explanation of the $\Delta I = 1/2$ rule through Q_2 penguin-like diagrams and a matrix element of the gluonic penguin Q_6 around three times the factorizable contribution.

After these results we can safely neglect the EM penguin contribution to $\Delta g_{C(N)}$ and conclude that these asymmetries are to a very good approximation proportional to $\text{Im } G_8$. This is interesting since with a good enough accuracy, this asymmetry can check a large

$\text{Im } G_8$. This makes also this asymmetry complementary to the direct CP-violating parameter ε'_K where there is a strong cancellation between the $\text{Im } G_8$ and $\text{Im } G_E$ contributions.

The results we get at LO are

$$\Delta g_C \Big|_{\text{LO}} = -[0.51 \pm 0.25] \times 10^{-4} \quad \text{and} \quad \Delta g_N \Big|_{\text{LO}} = [1.70 \pm 0.85] \times 10^{-5}, \quad (29)$$

where we have used the results in (24) and (28).

At NLO, we have to include several ingredients. For the real part, we need the chiral logs and counterterms, K_i . The chiral logs are analytically known^{12,1}. The real part of the counterterms, $\text{Re } K_i$, that enter in $K \rightarrow 3\pi$ can be obtained from the fit to data done in¹².

At NLO we also need $\text{Im } G'_8$ in addition to $\text{Im } G_8$, for which, to the best of our knowledge, there is just one calculation at NLO in $1/N_c$ at present in³¹. The results found there, using the same hadronic model discussed above, are

$$\text{Re } G'_8 = 0.9 \pm 0.1, \quad \text{and} \quad \text{Im } G'_8 = (1.0 \pm 0.2) \text{Im } \tau. \quad (30)$$

The imaginary part of the rest of the counterterms, $\text{Im } K_i$, is much more problematic and there is no NLO in $1/N_c$ calculation of them at present.

We can use several approaches to get the order of magnitude and/or the signs of the rest of the imaginary parts of the purely order p^4 couplings. These approaches to get information of the those unknown imaginary parts are factorization plus meson dominance³⁶, large N_c , or assuming that the ratio of the real and imaginary parts are dominated by the same dynamics at all orders in CHPT, therefore

$$\frac{\text{Im } K_i}{\text{Re } K_i} \simeq \frac{\text{Im } G_8}{\text{Re } G_8} \simeq \frac{\text{Im } G'_8}{\text{Re } G'_8}. \quad (31)$$

Notice that the last relation is well satisfied by the model calculation in³¹. This will allow us to check the counterterm dependence of the CP-violating asymmetries.

Using the estimates of the imaginary part of the counterterms above and the results from the fit in¹² for the real part, we have checked that the sum $g(K^+ \rightarrow \pi^+\pi^+\pi^-) + g(K^- \rightarrow \pi^-\pi^-\pi^+)$ is dominated by chiral logs. In the difference $g(K^+ \rightarrow \pi^+\pi^+\pi^-) - g(K^- \rightarrow \pi^-\pi^-\pi^+)$ the counterterm contribution is however important and needs more work.

The last ingredient to have a complete analysis of the CP-violating asymmetries at NLO are the phases generated by the two-loop amplitudes $K \rightarrow 3\pi$. The lowest non-trivial order is p^4 which is analytically known. This is the one that has to be used in LO prediction of Δg_C . At NLO, one needs the same calculation at order p^6 which is not available at present. However, one can do a very good job using the known results at order p^4 and the optical theorem to get analytically the order p^6 imaginary parts that come from two one-loop bubbles which we can assume to be dominant^{1,22} to a very good accuracy. We disregard three-body rescattering since they cannot be written as a bubble resummation. One can expect them to be rather small being suppressed by the phase space available.

We have presented here an update of the $K^+ \rightarrow 3\pi$ CP-violating asymmetries $\Delta g_{C(N)}$ at LO in the Standard Model. We are working in the full NLO corrections which will be ready soon¹.

The measurement of this decay by NA48 or elsewhere will be extremely interesting for several reasons. We have seen that the asymmetries $\Delta g_{C(N)}$ are proportional at lowest order to $\text{Im } G_8$ to a very good approximation. This allows a consistency check between the theoretical predictions for ε'_K ^{31,35,38}. Any prediction for ε'_K has to be also able to predict the CP-violating asymmetries. The measurement of those asymmetries may allow to check the large $\text{Im } G_8$ results found at NLO, see for instance^{31,35} and references therein.

Moreover, it seems that some models beyond the Standard Model can achieve values not much larger than 1×10^{-4} , see for instance³⁷. It will be important to check if that value can be attained in the Standard Model too.

We are also finishing the analysis of the CP-violating asymmetries in the integrated rates of the charged Kaon decays into three pions, $\Delta\Gamma$ at NLO order in CHPT.

Acknowledgments

This work has been supported in part by the RTN European Union No HPRN-CT-2002-00311 (EURIDICE). E.G. is indebted to MECD (Spain) for a F.P.U. fellowship. The work of E.G. and J.P. has been supported in part by MCYT (Spain) under Grants No. FPA2000-1558 and HF2001-0116 and by Junta de Andalucía Grant No. FQM-101. The work of I.S. has been supported in part by the Swiss National Science Foundation and RTN, BBW-Contract No. 01.0357.

References

1. E. Gámiz, J. Prades, and I. Scimemi, preprint CAFPE/19-03, Granada preprint UGFT/149-03, and Bern preprint BUTP-2003-12.
2. S. Weinberg, *Physica A* **96** (1979) 327.
3. J. Gasser and H. Leutwyler, *Annals Phys.* **158** (1984) 142.
4. J. Gasser and H. Leutwyler, *Nucl. Phys. B* **250** (1985) 465.
5. G. Ecker, Lectures given at the Advanced School on QCD 2000, Benasque, (Huesca), Spain, hep-ph/0011026; A. Pich, hep-ph/9806303
6. J. Kambor, J. Missimer and D. Wyler, *Nucl. Phys. B* **346** (1990) 17.
7. G. Esposito-Farèse *Z. Phys. C* **50** (1991) 255.
8. G. Ecker, J. Kambor and D. Wyler, *Nucl. Phys. B* **394** (1993) 101.
9. G. Ecker, G. Isidori, G. Müller, H. Neufeld and A. Pich, *Nucl. Phys. B* **591** (2000) 419.
10. B.R. Holstein, *Phys. Rev.* **177** (1969) 2417; *ibidem* **183** (1969) 1228.
11. J. Kambor, J. Missimer and D. Wyler, *Phys. Lett. B* **261** (1991) 496.
12. J. Bijnens, P. Dhonte and F. Persson, *Nucl. Phys. B* **648** (2003) 317.
13. J. Bijnens, E. Pallante and J. Prades, *Nucl. Phys. B* **521** (1998) 305.
14. L.-F. Li and L. Wolfenstein, *Phys. Rev. D* **21** (1980) 178.
15. C. Avilez, *Phys. Rev. D* **23** (1981) 1124.
16. B. Grinstein, S.-J. Rey and M.B. Wise, *Phys. Rev. D* **33** (1986) 1495.
17. J.F. Donoghue, B.R. Holstein and G. Valencia, *Phys. Rev. D* **36** (1987) 798.
18. A.A. Bel'kov, A.V. Lanyov, G. Bohm and D. Ebert, *Phys. Lett. B* **232** (1989) 118; A.A. Bel'kov, G. Bohm, D. Ebert, A.V. Lanyov and A. Schaale, *Int. J. Mod. Phys. A* **7** (1992) 4757.
19. G. D'Ambrosio, G. Isidori and N. Paver, *Phys. Lett. B* **273** (1991) 497.
20. G. Isidori, L. Maiani and A. Pugliese, *Nucl. Phys. B* **381** (1992) 522.
21. E.P. Shabalin, *Nucl. Phys. B* **409** (1993) 87; ITEP-92-6; *Phys. Atom. Nucl.* **61** (1998) 1372 [*Yad. Fiz.* **61** (1998) 1478].
22. G. D'Ambrosio, G. Isidori, A. Pugliese and N. Paver, *Phys. Rev. D* **50** (1994) 5767 [Erratum-*ibid.* **51** (1995) 3975].
23. R. Wanke, Talk at the XXXVIII Rencontres de Moriond on ElectroWeak Interactions and Unified Theories, Les Arcs, France, March 15-22 2003
24. J. Bijnens, J. Prades and E. de Rafael, *Phys. Lett. B* **348** (1995) 226.
25. V. Cirigliano, J.F. Donoghue, E. Golowich and K. Maltman, *Phys. Lett. B* **555** (2003) 71; hep-ph/0209332; *Phys. Lett. B* **522** (2001) 245.
26. S. Narison, *Nucl. Phys. B* **593** (2001) 3.
27. J. Bijnens, E. Gámiz and J. Prades, *J. High Energy Phys.* **10** (2001) 009; E. Gámiz, J. Prades and J. Bijnens, hep-ph/0209089.
28. M. Knecht, S. Peris and E. de Rafael, *Phys. Lett. B* **508** (2001) 117; *Phys. Lett. B* **457** (1999) 227.
29. J.I. Noaki *et al.* [CP-PACS Collaboration], hep-lat/0108013; T. Blum *et al.* [RBC Collaboration], hep-lat/0110075;
30. D. Becirevic *et al.* [SPQCDR Collaboration], hep-lat/0209136.
31. J. Bijnens and J. Prades, *J. High Energy Phys.* **06**, 035 (2000); *Nucl. Phys. Proc. Suppl.* **96**, 354 (2001); hep-ph/0009156; hep-ph/0009155;
32. J. Bijnens and J. Prades, *J. High Energy Phys.* **01**, 023 (1999); J. Prades, *Nucl. Phys. Proc. Suppl.* **86**, 294 (2000).
33. S. Peris, M. Perrottet and E. de Rafael, *J. High Energy Phys.* **05** (1998) 011.

- 34. J. Bijnens, E. Gámiz, E. Lipartia and J. Prades, J. High Energy Phys. **04** (2003) 055.
- 35. T. Hambye, S. Peris and E. de Rafael, hep-ph/0305104
- 36. G. Ecker, J. Gasser, A. Pich and E. de Rafael, Nucl. Phys. B **321** (1989) 311.
- 37. G. D'Ambrosio, G. Isidori and G. Martinelli, Phys. Lett. B **480** (2000) 164.
- 38. E. Pallante, A. Pich and I. Scimemi, Nucl. Phys. B **617** (2001) 441; E. Pallante and A. Pich, Nucl. Phys. B **592** (2001) 294; Phys. Rev. Lett. **84** (2000) 2568.